

INFORMATION FOR THE PATIENT

Brief Summary Patient Package Insert

Triphasil[®]-28

Tablets

(levonorgestrel and ethinyl estradiol tablets—triphasic regimen)

R_x only

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

28 tablets containing the following:

6 brown tablets—0.050 mg levonorgestrel + 0.030 mg ethinyl estradiol;

5 white tablets—0.075 mg levonorgestrel + 0.040 mg ethinyl estradiol;

10 light-yellow tablets—0.125 mg levonorgestrel + 0.030 mg ethinyl estradiol;
plus 7 light-green tablets—inert.

Oral contraceptives, also known as “birth-control pills” or “the pill,” are taken to prevent pregnancy, and when taken correctly, have a failure rate of less than 1.0% per year when used without missing any pills. The average failure rate of large numbers of pill users is 5% per year when women who miss pills are included. For most women oral contraceptives are also free of serious or unpleasant side effects. However, forgetting to take pills considerably increases the chances of pregnancy.

For the majority of women, oral contraceptives can be taken safely. But there are some women who are at high risk of developing certain serious diseases that can be life-threatening or may cause temporary or permanent disability or death. The risks associated with taking oral contraceptives increase significantly if you:

- smoke.
- have high blood pressure, diabetes, high cholesterol, or a tendency to form blood clots, or are obese.
- have or have had clotting disorders, heart attack, stroke, angina pectoris, cancer of the breast or sex organs, jaundice, or malignant or benign liver tumors.

You should not take the pill if you suspect you are pregnant or have unexplained vaginal bleeding.

Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels from oral-contraceptive use. This risk increases with age and with the amount of smoking (15 or more cigarettes per day has been associated with a significantly increased risk) and is quite marked in women over 35 years of age. Women who use oral contraceptives should not smoke.

Most side effects of the pill are not serious. The most common such effects are nausea, vomiting, bleeding between menstrual periods, weight gain, breast tenderness, and difficulty wearing contact lenses. These side effects, especially nausea and vomiting, may subside within the first three months of use.

The serious side effects of the pill occur very infrequently, especially if you are in good health and do not smoke. However, you should know that the following medical conditions have been associated with or made worse by the pill:

1. Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), stoppage or rupture of a blood vessel in the brain (stroke), blockage of blood vessels in the heart (heart attack and angina pectoris) or other organs of the body. As mentioned above, smoking increases the risk of heart attacks and strokes and subsequent serious medical consequences. Women with migraine also may be at increased risk of stroke.
2. Liver tumors, which may rupture and cause severe bleeding. A possible but not definite association has been found with the pill and liver cancer. However, liver cancers are extremely rare. The chance of developing liver cancer from using the pill is thus even rarer.
3. High blood pressure, although blood pressure usually returns to normal when the pill is stopped.

The symptoms associated with these serious side effects are discussed in the detailed leaflet given to you with your supply of pills. Notify your doctor or health-care provider if you notice any unusual physical disturbances while taking the pill. In addition, drugs such as rifampin, as well as some anticonvulsants and some antibiotics, and possibly St. John's wort, may decrease oral-contraceptive effectiveness.

Breast cancer has been diagnosed slightly more often in women who use the pill than in women of the same age who do not use the pill. This very small increase in the number of breast cancer diagnoses gradually disappears during the 10 years after stopping use of the pill. It is not known whether the difference is caused by the pill. It may be that women taking the pill were examined more often, so that breast cancer was more likely to be detected.

Some studies have found an increase in the incidence of cancer or precancerous lesions of the cervix in women who use the pill. However, this finding may be related to factors other than the use of the pill.

Taking the pill provides some important noncontraceptive benefits. These include less painful menstruation, less menstrual blood loss and anemia, fewer pelvic infections, and fewer cancers of the ovary and the lining of the uterus.

Be sure to discuss any medical condition you may have with your health-care provider. Your health-care provider will take a medical and family history before prescribing oral contraceptives and will examine you. The physical examination may be delayed to another time if you request it and the health-care provider believes that it is appropriate to postpone it. You should be reexamined at least once a year while taking oral contraceptives. The detailed patient information leaflet gives you further information which you should read and discuss with your health-care provider.

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against transmission of HIV (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

Instructions for the Patient


HOW TO USE THE TRIPHASIL®-28 MINI-PACK™

IMPORTANT: Do not turn the clear outer ring of the dispenser before you read these instructions. The clear ring has been preset to your first tablet.

STEP 1—Set the white inner dial of your first pack to your starting day by using a coin in the center slot.

Day 1 Start: Turn the white dial until the day that corresponds to your first day of bleeding lines up with the starting window. **FOR EXAMPLE:** For your first pack, if your first day of bleeding falls on a Wednesday, turn the white dial until “WED” appears under the starting window. (The starting window in the clear ring is below the “U” notch.)

Sunday Start: Turn the white dial to “SUN.”




STEP 2—To open the starting window, **push up** on the tab to break it off and begin pill taking where it reads “START HERE.”

Day 1 Start: Take the first brown “active” pill on the first day of your period. You do not need to use a back-up birth-control method since you are starting the pill at the beginning of your period.

Sunday Start: Take the first brown “active” pill of the first pack on the *Sunday after your period starts*, even if you are still bleeding. If your period begins on Sunday, start the pack that same day. *Use another method of birth control* as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days).

STEP 3—Take 1 pill each day by turning the clear ring in the direction of the arrow. After you have taken the last light-yellow pill, you will usually have a period a few days later, which will be during the time when you are taking the light-green pills. Don’t be alarmed if the amount of bleeding is not the same as before. Take a pill every day without interruption until the last light-green pill has been taken.

STEP 4—After taking the last light-green pill, start a new dial dispenser the **very next day** no matter when your period started. Remember to set the white dial of your new pack to your starting day. If you have taken the pills as directed, the first brown pill in every new dispenser will always be taken on the same day of the week that you first began taking Triphasil[®] (levonorgestrel and ethinyl estradiol tablets—triphasic regimen). If you choose the **Sunday Start** regimen, this will always be on Sunday. Any time you start the pills later than directed, protect yourself by using another method of birth control until you have taken a pill a day for seven consecutive days (6 brown and 1 white).



WHAT TO DO WHEN YOU SWITCH YOUR BRAND OF PILL

Start the next pack on the day after your last “reminder” pill. Do not wait any days between packs.

WHAT TO DO IF YOU HAVE SPOTTING OR BLEEDING

If spotting or bleeding should occur at an unexpected time while on the pills, continue to take them as directed. This is usually temporary and without significance; however, if bleeding is persistent or prolonged, consult your physician. Also call your doctor about any problem which may bother you.

WHAT TO DO IF YOU HAVE VOMITING

*IF YOU HAVE VOMITING (within 3 to 4 hours after you take your pill), you should follow the instructions for **WHAT TO DO IF YOU MISS PILLS**. IF YOU HAVE DIARRHEA or IF YOU TAKE SOME MEDICINES, including some antibiotics, your pills may not work as well. Use a back-up method (such as condoms, spermicide, or sponge) until you check with your doctor or clinic.*

WHAT TO DO IF YOU MISS A PERIOD

In rare instances a period may be skipped. If you have taken pills as directed, continue as before. However, if you did not take your pills as directed (missed one or more pills or started taking them on a day later than you should have) and miss a period, or if you miss two successive periods even though you followed directions, stop taking your pills and call your physician promptly since pregnancy should be ruled out before you continue medication. Another method of birth control other than oral contraceptives should be used during this time.

WHAT TO DO IF YOU MISS PILLS

The pill may not be as effective if you miss brown, white or light-yellow “active” pills, and particularly if you miss the first few brown or the last few light-yellow “active” pills in a pack.

If you **MISS 1** brown, white or light-yellow “active” pill:

1. Take it as soon as you remember. Take the next pill at your regular time. This means you may take 2 pills in 1 day.
2. You do not need to use a back-up birth-control method if you have sex.

If you **MISS 2** brown, white or light-yellow “active” pills in a row in **WEEK 1 OR WEEK 2** of your pack:

1. Take 2 pills on the day you remember and 2 pills the next day.
2. Then take 1 pill a day until you finish the pack.
3. You **MAY BECOME PREGNANT** if you have sex in the *7 days* after you miss pills. You **MUST** use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

If you **MISS 2** brown, white or light-yellow “active” pills in a row in **THE 3rd WEEK:**

1. ***If you are a Day 1 Starter:***

THROW OUT the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your doctor or clinic because you might be pregnant.

3. You MAY BECOME PREGNANT if you have sex in the *7 days* after you miss pills. You MUST use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

If you **MISS 3 OR MORE** brown, white or light-yellow “active” pills in a row (during the first 3 weeks):

1. ***If you are a Day 1 Starter:***

THROW OUT the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your doctor or clinic because you might be pregnant.

3. You MAY BECOME PREGNANT if you have sex in the *7 days* after you miss pills. You MUST use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

For additional information see “[Detailed Patient Labeling.](#)”



This product's label may have been updated. For current package insert and further product information, please visit www.wyeth.com or call our medical communications department toll-free at 1-800-934-5556.



Wyeth[®]

Wyeth Pharmaceuticals Inc.
Philadelphia, PA 19101

W10507P002
ET01
Rev 06/06

DETAILED PATIENT LABELING

Triphasil[®] -21 & -28

Tablets

(levonorgestrel and ethinyl estradiol tablets—triphasic regimen)

R_x only

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

INTRODUCTION

Any woman who considers using oral contraceptives (the birth-control pill or the pill) should understand the benefits and risks of using this form of birth control. This leaflet will give you much of the information you will need to make this decision and will also help you determine if you are at risk of developing any of the serious side effects of the pill. It will tell you how to use the pill properly so that it will be as effective as possible. However, this leaflet is not a replacement for a careful discussion between you and your health-care provider. You should discuss the information provided in this leaflet with him or her, both when you first start taking the pill and during your revisits. You should also follow your health-care provider's advice with regard to regular check-ups while you are on the pill.

EFFECTIVENESS OF ORAL CONTRACEPTIVES

Oral contraceptives or “birth-control pills” or “the pill” are used to prevent pregnancy and are more effective than other nonsurgical methods of birth control. When they are taken correctly, the chance of becoming pregnant is less than 1.0% when used perfectly, without missing any pills. Average failure rates are 5% per year. The chance of becoming pregnant increases with each missed pill during the menstrual cycle.

In comparison, average failure rates for other nonsurgical methods of birth control during the first year of use are as follows:

TABLE: PERCENTAGE OF WOMEN EXPERIENCING AN UNINTENDED PREGNANCY DURING THE FIRST YEAR OF USE OF A CONTRACEPTIVE METHOD

Method	Perfect Use	Average Use
Levonorgestrel implants	0.05	0.05
Male sterilization	0.1	0.15
Female sterilization	0.5	0.5
Depo-Provera [®] (injectable progestogen)	0.3	0.3
Oral contraceptives		5
Combined	0.1	NA
Progestin only	0.5	NA

Method	Perfect Use	Average Use
IUD		
Progesterone	1.5	2.0
Copper T 380A	0.6	0.8
Condom (male) without spermicide	3	14
(female) without spermicide	5	21
Cervical cap		
Never given birth	9	20
Given birth	26	40
Vaginal Sponge		
Never given birth	9	20
Given birth	20	40
Diaphragm with spermicidal cream or jelly	6	20
Spermicides alone (foam, creams, jellies, and vaginal suppositories)	6	26
Periodic abstinence (all methods)	1-9*	25
Withdrawal	4	19
No contraception (planned pregnancy)	85	85

NA – not available

*Depending on method (calendar, ovulation, symptothermal, post-ovulation)

Adapted from Hatcher RA et al, *Contraceptive Technology: 17th Revised Edition*. NY, NY: Ardent Media, Inc., 1998.

WHO SHOULD NOT TAKE ORAL CONTRACEPTIVES

Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels from oral-contraceptive use. This risk increases with age and with the amount of smoking (15 or more cigarettes per day has been associated with a significantly increased risk) and is quite marked in women over 35 years of age. Women who use oral contraceptives should not smoke.

Some women should not use the pill. For example, you should not take the pill if you are pregnant or think you may be pregnant. You should also not use the pill if you have had any of the following conditions:

- Heart attack or stroke.
- Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), or eyes.
- Blood clots in the deep veins of your legs.
- Known or suspected breast cancer or cancer of the lining of the uterus, cervix, or vagina or certain hormonally-sensitive cancers.
- Liver tumor (benign or cancerous).

Or, if you have any of the following:

- Chest pain (angina pectoris).
- Unexplained vaginal bleeding (until a diagnosis is reached by your doctor).
- Yellowing of the whites of the eyes or of the skin (jaundice) during pregnancy or during previous use of the pill.
- Known or suspected pregnancy.
- Heart valve or heart rhythm disorders that may be associated with formation of blood clots.
- Diabetes affecting your circulation.
- Uncontrolled high blood pressure.
- Active liver disease with abnormal liver function tests.
- Allergy or hypersensitivity to any of the components of Triphasil[®] (levonorgestrel and ethinyl estradiol tablets – triphasic regimen).

Tell your health-care provider if you have ever had any of these conditions. Your health-care provider can recommend another method of birth control.

OTHER CONSIDERATIONS BEFORE TAKING ORAL CONTRACEPTIVES

Tell your health-care provider if you or any family member has ever had:

- Breast nodules, fibrocystic disease of the breast, an abnormal breast X-ray or mammogram.
- Diabetes.
- Elevated cholesterol or triglycerides.
- High blood pressure.
- A tendency to form blood clots.
- Migraine or other headaches or epilepsy.
- Mental depression.
- Gallbladder, heart, or kidney disease.
- History of scanty or irregular menstrual periods.

Women with any of these conditions should be checked often by their health-care provider if they choose to use oral contraceptives. Also, be sure to inform your doctor or health-care provider if you smoke or are on any medications.

RISKS OF TAKING ORAL CONTRACEPTIVES

1. Risk of developing blood clots

Blood clots and blockage of blood vessels are the most serious side effects of taking oral contraceptives and can be fatal. In particular, a clot in the legs can cause thrombophlebitis and a clot that travels to the lungs can cause a sudden blocking of the vessel carrying blood to the lungs. Rarely, clots occur in the blood vessels of the eye and may cause blindness, double vision, or impaired vision.

If you take oral contraceptives and need elective surgery, need to stay in bed for a prolonged illness, or have recently delivered a baby, you may be at risk of developing blood clots. You should consult your doctor about stopping oral contraceptives three to four weeks before surgery and not taking oral contraceptives for two weeks after surgery or during bed rest. You should also not take oral contraceptives soon after delivery of a baby or a midtrimester pregnancy termination. It is advisable to wait for at least four weeks after delivery if you are not breast-feeding. If you are breast-feeding, you should wait until you have weaned your child before using the pill. (See also the section on breast-feeding in “[GENERAL PRECAUTIONS](#)”.)

2. Heart attacks and strokes

Oral contraceptives may increase the tendency to develop strokes (stoppage or rupture of blood vessels in the brain) and angina pectoris and heart attacks (blockage of blood vessels in the heart). Any of these conditions can cause death or serious disability.

Smoking greatly increases the possibility of suffering heart attacks and strokes. Furthermore, smoking and the use of oral contraceptives greatly increase the chances of developing and dying of heart disease.

Women with migraine (especially migraine with aura) who take oral contraceptives also may be at higher risk of stroke.

3. Gallbladder disease

Oral-contraceptive users probably have a greater risk than nonusers of having gallbladder disease, although this risk may be related to pills containing high doses of estrogens.

4. Liver tumors

In rare cases, oral contraceptives can cause benign but dangerous liver tumors. These benign liver tumors can rupture and cause fatal internal bleeding. In addition, a possible but not definite association has been found with the pill and liver cancers in two studies in which a few women who developed these very rare cancers were found to have used oral contraceptives for long periods. However, liver cancers are extremely rare. The chance of developing liver cancer from using the pill is thus even rarer.

5. Cancer of the reproductive organs

Breast cancer has been diagnosed slightly more often in women who use the pill than in women of the same age who do not use the pill. This very small increase in the number of breast cancer diagnoses gradually disappears during the 10 years after stopping use of the pill. It is not known whether the difference is caused by the pill. It may be that women taking the pill were examined more often, so that breast cancer was more likely to be detected.

Some studies have found an increase in the incidence of cancer or precancerous lesions of the cervix in women who use oral contraceptives. However, this finding may be related to factors other than the use of oral contraceptives.

6. *Lipid metabolism and inflammation of the pancreas*

In patients with inherited defects of lipid metabolism, there have been reports of significant elevations of plasma triglycerides during estrogen therapy. This has led to pancreatitis in some cases.

ESTIMATED RISK OF DEATH FROM A BIRTH-CONTROL METHOD OR PREGNANCY

All methods of birth control and pregnancy are associated with a risk of developing certain diseases which may lead to disability or death. An estimate of the number of deaths associated with different methods of birth control and pregnancy has been calculated and is shown in the following [table](#).

ANNUAL NUMBER OF BIRTH-RELATED OR METHOD-RELATED DEATHS
ASSOCIATED WITH CONTROL OF FERTILITY PER 100,000 NONSTERILE WOMEN, BY
FERTILITY-CONTROL METHOD ACCORDING TO AGE

Method of control and outcome	15-19	20-24	25-29	30-34	35-39	40-44
No fertility control methods*	7.0	7.4	9.1	14.8	25.7	28.2
Oral contraceptives nonsmoker**	0.3	0.5	0.9	1.9	13.8	31.6
Oral contraceptives smoker**	2.2	3.4	6.6	13.5	51.1	117.2
IUD**	0.8	0.8	1.0	1.0	1.4	1.4
Condom*	1.1	1.6	0.7	0.2	0.3	0.4
Diaphragm/ spermicide*	1.9	1.2	1.2	1.3	2.2	2.8
Periodic abstinence*	2.5	1.6	1.6	1.7	2.9	3.6

*Deaths are birth related

**Deaths are method related

In the above table, the risk of death from any birth-control method is less than the risk of childbirth, except for oral-contraceptive users over the age of 35 who smoke and pill users over the age of 40 even if they do not smoke. It can be seen in the table that for women aged 15 to 39, the risk of death was highest with pregnancy (7 to 26 deaths per 100,000 women, depending on age). Among pill users who do not smoke, the risk of death was always lower than that associated with pregnancy for any age group, except for those women over the age of 40, when the risk increases to 32 deaths per 100,000 women, compared to 28 associated with pregnancy at that age. However, for pill users who smoke and are over the age of 35, the estimated number of deaths exceeds those for other methods of birth control. If a woman is over the age of 40 and smokes, her estimated risk of death is four times higher (117/100,000 women) than the estimated risk associated with pregnancy (28/100,000 women) in that age group.

The suggestion that women over 40 who don't smoke should not take oral contraceptives is based on information from older high-dose pills and on less-selective use of pills than is practiced today. An Advisory Committee of the FDA discussed this issue in 1989 and recommended that the benefits of oral-contraceptive use by healthy, nonsmoking women over 40 years of age may outweigh the possible risks. However, all women, especially older women, are cautioned to use the lowest-dose pill that is effective.

WARNING SIGNALS

If any of these adverse effects occur while you are taking oral contraceptives, call your doctor immediately:

- Sharp chest pain, coughing of blood, or sudden shortness of breath (indicating a possible clot in the lung).
- Pain in the calf (indicating a possible clot in the leg).
- Crushing chest pain or heaviness in the chest (indicating a possible heart attack).
- Sudden severe headache or vomiting, dizziness or fainting, disturbances of vision or speech, weakness, or numbness in an arm or leg (indicating a possible stroke).
- Sudden partial or complete loss of vision (indicating a possible clot in the eye).
- Breast lumps (indicating possible breast cancer or fibrocystic disease of the breast; ask your doctor or health-care provider to show you how to examine your breasts).
- Severe pain or tenderness in the stomach area (indicating a possibly ruptured liver tumor).
- Difficulty in sleeping, weakness, lack of energy, fatigue, or change in mood (possibly indicating severe depression).
- Jaundice or a yellowing of the skin or eyeballs, accompanied frequently by fever, fatigue, loss of appetite, dark-colored urine, or light-colored bowel movements (indicating possible liver problems).

SIDE EFFECTS OF ORAL CONTRACEPTIVES

1. *Vaginal bleeding*

Irregular vaginal bleeding or spotting may occur while you are taking the pills. Irregular bleeding may vary from slight staining between menstrual periods to breakthrough bleeding which is a flow much like a regular period. Irregular bleeding occurs most often during the first few months of oral-contraceptive use, but may also occur after you have been taking the pill for some time. Such bleeding may be temporary and usually does not indicate any serious problems. It is important to continue taking your pills on schedule. If the bleeding occurs in more than one cycle or lasts for more than a few days, talk to your doctor or health-care provider.

2. *Contact lenses*

If you wear contact lenses and notice a change in vision or an inability to wear your lenses, contact your doctor or health-care provider.

3. *Fluid retention*

Oral contraceptives may cause edema (fluid retention) with swelling of the fingers or ankles and may raise your blood pressure. If you experience fluid retention, contact your doctor or health-care provider.

4. *Melasma*

A spotty darkening of the skin is possible, particularly of the face.

5. *Other side effects*

Other side effects may include nausea, breast tenderness, change in appetite, headache, nervousness, depression, dizziness, loss of scalp hair, rash, vaginal infections, inflammation of the pancreas, and allergic reactions.

If any of these side effects bother you, call your doctor or health-care provider.

GENERAL PRECAUTIONS

1. *Missed periods and use of oral contraceptives before or during early pregnancy*

There may be times when you may not menstruate regularly after you have completed taking a cycle of pills. If you have taken your pills regularly and miss one menstrual period, continue taking your pills for the next cycle but be sure to inform your health-care provider before doing so. If you have not taken the pills daily as instructed and missed a menstrual period, or if you missed two consecutive menstrual periods, you may be pregnant. Check with your health-care provider immediately to determine whether you are pregnant. Do not continue to take oral contraceptives until you are sure you are not pregnant, but continue to use another method of contraception.

There is no conclusive evidence that oral-contraceptive use is associated with an increase in birth defects when taken inadvertently during early pregnancy. Previously, a few studies had reported that oral contraceptives might be associated with birth defects, but these studies have not been confirmed. Nevertheless, oral contraceptives or any other drugs should not be used during pregnancy unless clearly necessary and prescribed by your doctor. You should check with your doctor about risks to your unborn child of any medication taken during pregnancy.

2. *While breast-feeding*

If you are breast-feeding, consult your doctor before starting oral contraceptives. Some of the drug will be passed on to the child in the milk. A few adverse effects on the child have been reported, including yellowing of the skin (jaundice) and breast enlargement. In addition, oral contraceptives may decrease the amount and quality of your milk. If possible, do not use oral contraceptives while breast-feeding. You should use another method of contraception since breast-feeding provides only partial protection from becoming pregnant, and this partial protection decreases significantly as you breast-feed for longer periods of time. You should consider starting oral contraceptives only after you have weaned your child completely.

3. *Laboratory tests*

If you are scheduled for any laboratory tests, tell your doctor you are taking birth-control pills. Certain blood tests may be affected by birth-control pills.

4. *Drug interactions*

Certain drugs may interact with birth-control pills to make them less effective in preventing pregnancy or cause an increase in breakthrough bleeding. Such drugs include rifampin, drugs used for epilepsy such as barbiturates (for example, phenobarbital) and phenytoin (Dilantin[®] is one brand of this drug), primidone (Mysoline[®]), topiramate (Topamax[®]), phenylbutazone (Butazolidin[®] is one brand), some drugs used for HIV such as ritonavir (Norvir[®]), modafinil (Provigil[®]), and possibly certain antibiotics (such as ampicillin and other penicillins, and tetracyclines) and St. John's wort. You may need to use an additional method of contraception during any cycle in which you take drugs that can make oral contraceptives less effective.

You may be at higher risk of a specific type of liver dysfunction if you take troleandomycin and oral contraceptives at the same time.

You should inform your healthcare provider about all medicines you are taking, including nonprescription products.

5. *Sexually transmitted diseases*

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against transmission of HIV (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

HOW TO TAKE THE PILL

IMPORTANT POINTS TO REMEMBER

BEFORE YOU START TAKING YOUR PILLS:

1. BE SURE TO READ THESE DIRECTIONS:

Before you start taking your pills.
Anytime you are not sure what to do.

2. THE RIGHT WAY TO TAKE THE PILL IS TO TAKE ONE PILL EVERY DAY AT THE SAME TIME.

If you miss pills you could get pregnant. This includes starting the pack late. The more pills you miss, the more likely you are to get pregnant.

3. MANY WOMEN HAVE SPOTTING OR LIGHT BLEEDING, OR MAY FEEL SICK TO THEIR STOMACH DURING THE FIRST 1-3 PACKS OF PILLS.

If you feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it doesn't go away, check with your doctor or clinic.

4. MISSING PILLS CAN ALSO CAUSE SPOTTING OR LIGHT BLEEDING, even when you make up these missed pills.

On the days you take 2 pills to make up for missed pills, you could also feel a little sick to your stomach.

*5. IF YOU HAVE VOMITING (within 3 to 4 hours after you take your pill), you should follow the instructions for **WHAT TO DO IF YOU MISS PILLS**. IF YOU HAVE DIARRHEA, or IF YOU TAKE SOME MEDICINES, including some antibiotics, your pills may not work as well.*

Use a back-up method (such as condoms, spermicide, or sponge) until you check with your doctor or clinic.

6. IF YOU HAVE TROUBLE REMEMBERING TO TAKE THE PILL, talk to your doctor or clinic about how to make pill-taking easier or about using another method of birth control.

7. IF YOU HAVE ANY QUESTIONS OR ARE UNSURE ABOUT THE INFORMATION IN THIS LEAFLET, call your doctor or clinic.

BEFORE YOU START TAKING YOUR PILLS

1. DECIDE WHAT TIME OF DAY YOU WANT TO TAKE YOUR PILL.

It is important to take it at about the same time every day.


2. LOOK AT YOUR PILL PACK TO SEE IF IT HAS 21 OR 28 PILLS:

The *21-pill pack* has 21 “active” brown, white or light-yellow pills (with hormones) to take for 3 weeks, followed by 1 week without pills.

The *28-pill pack* has 21 “active” brown, white or light-yellow pills (with hormones) to take for 3 weeks, followed by 1 week of reminder light-green pills (without hormones).

3. ALSO FIND:

- 1) where on the pack to start taking pills.
- 2) in what order to take the pills (follow the arrow).



4. BE SURE YOU HAVE READY AT ALL TIMES:

ANOTHER KIND OF BIRTH CONTROL (such as condoms, spermicide, or sponge) to use as a back-up in case you miss pills.

AN EXTRA, FULL PILL PACK.

WHEN TO START THE FIRST PACK OF PILLS

You have a choice of which day to start taking your first pack of pills. Decide with your doctor or clinic which is the best day for you. Pick a time of day which will be easy to remember.

DAY 1 START:

1. Take the first “active” brown pill of the first pack during the *first 24 hours of your period*.
2. You will not need to use a back-up method of birth control, since you are starting the pill at the beginning of your period.

SUNDAY START:

1. Take the first “active” brown pill of the first pack on the *Sunday after your period starts*, even if you are still bleeding. If your period begins on Sunday, start the pack that same day.
2. *Use another method of birth control* as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days). Condoms, spermicide, or the sponge are good back-up methods of birth control.

WHAT TO DO DURING THE MONTH

1. TAKE ONE PILL AT THE SAME TIME EVERY DAY UNTIL THE PACK IS EMPTY.

Do not skip pills even if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea).

Do not skip pills even if you do not have sex very often.

2. WHEN YOU FINISH A PACK OR SWITCH YOUR BRAND OF PILLS:

21 pills: Wait 7 days to start the next pack. You will probably have your period during that week. Be sure that no more than 7 days pass between 21-day packs.

28 pills: Start the next pack on the day after your last “reminder” pill. Do not wait any days between packs.

WHAT TO DO IF YOU MISS PILLS

The pill may not be as effective if you miss brown, white or light-yellow “active” pills, and particularly if you miss the first few brown or the last few light-yellow “active” pills in a pack.

If you **MISS 1** brown, white or light-yellow “active” pill:

1. Take it as soon as you remember. Take the next pill at your regular time. This means you may take 2 pills in 1 day.
2. You do not need to use a back-up birth-control method if you have sex.

If you **MISS 2** brown, white or light-yellow “active” pills in a row in **WEEK 1 OR WEEK 2** of your pack:

1. Take 2 pills on the day you remember and 2 pills the next day.
2. Then take 1 pill a day until you finish the pack.
3. You **MAY BECOME PREGNANT** if you have sex in the *7 days* after you miss pills. You **MUST** use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

If you **MISS 2** brown, white or light-yellow “active” pills in a row in **THE 3rd WEEK**:

1. ***If you are a Day 1 Starter:***

THROW OUT the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday.

On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your doctor or clinic because you might be pregnant.
3. You **MAY BECOME PREGNANT** if you have sex in the *7 days* after you miss pills. You **MUST** use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

If you **MISS 3 OR MORE** brown, white or light-yellow “active” pills in a row (during the first 3 weeks):

1. ***If you are a Day 1 Starter:***

THROW OUT the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday.

On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your doctor or clinic because you might be pregnant.
3. You **MAY BECOME PREGNANT** if you have sex in the *7 days* after you miss pills. You **MUST** use another birth-control method (such as condoms, spermicide, or sponge) as a back-up for those 7 days.

A REMINDER FOR THOSE ON 28-DAY PACKS

If you forget any of the 7 light-green “reminder” pills in Week 4:

THROW AWAY the pills you missed.

Keep taking 1 pill each day until the pack is empty.

You do not need a back-up method if you start your next pack on time.

FINALLY, IF YOU ARE STILL NOT SURE WHAT TO DO ABOUT THE PILLS YOU HAVE MISSED

Use a BACK-UP METHOD anytime you have sex.

KEEP TAKING ONE PILL EACH DAY until you can reach your doctor or clinic.

Pregnancy due to pill failure

The incidence of pill failure resulting in pregnancy is approximately less than 1.0% if taken every day as directed, but average failure rates are 5%. If you do become pregnant, the risk to the fetus is minimal, but you should stop taking your pills and discuss the pregnancy with your doctor.

Pregnancy after stopping the pill

There may be some delay in becoming pregnant after you stop using oral contraceptives, especially if you had irregular menstrual cycles before you used oral contraceptives. It may be advisable to postpone conception until you begin menstruating regularly once you have stopped taking the pill and desire pregnancy.

There does not appear to be any increase in birth defects in newborn babies when pregnancy occurs soon after stopping the pill.

Overdosage

Serious ill effects have not been reported following ingestion of large doses of oral contraceptives by young children. Overdosage may cause nausea and withdrawal bleeding in females. In case of overdosage, contact your health-care provider or pharmacist.

Other information

Your health-care provider will take a medical and family history before prescribing oral contraceptives and will examine you. The physical examination may be delayed to another time if you request it and the health-care provider believes that it is appropriate to postpone it. You should be reexamined at least once a year. Be sure to inform your health-care provider if there is a family history of any of the conditions listed previously in this leaflet. Be sure to keep all appointments with your health-care provider, because this is a time to determine if there are early signs of side effects of oral-contraceptive use.

Do not use the drug for any condition other than the one for which it was prescribed. This drug has been prescribed specifically for you; do not give it to others who may want birth-control pills.

HEALTH BENEFITS FROM ORAL CONTRACEPTIVES

In addition to preventing pregnancy, use of oral contraceptives may provide certain benefits. They are:

- Menstrual cycles may become more regular.
- Blood flow during menstruation may be lighter, and less iron may be lost. Therefore, anemia due to iron deficiency is less likely to occur.
- Pain or other symptoms during menstruation may be encountered less frequently.
- Ovarian cysts may occur less frequently.
- Ectopic (tubal) pregnancy may occur less frequently.
- Noncancerous cysts or lumps in the breast may occur less frequently.
- Acute pelvic inflammatory disease may occur less frequently.
- Oral-contraceptive use may provide some protection against developing two forms of cancer: cancer of the ovaries and cancer of the lining of the uterus.

If you want more information about birth-control pills, ask your doctor or pharmacist. They have a more technical leaflet called the Professional Labeling which you may wish to read.



This product's label may have been updated. For current package insert and further product information, please visit www.wyeth.com or call our medical communications department toll-free at 1-800-934-5556.



Wyeth[®]

Wyeth Pharmaceuticals Inc.
Philadelphia, PA 19101

W10507P002
ET01
Rev 06/06