

PRISTIQ Progress Tracker



The PRISTIQ Progress Tracker is a written record that will help you and your health care professional to monitor your progress while you are taking PRISTIQ. Fill out the Progress Tracker each day starting on Sunday and ending the following Saturday, completing one Progress Tracker every week for 8 weeks. Don't worry if you miss a day or two.

Just resume filling out the Progress Tracker the following day.

How to Use the PRISTIQ Progress Tracker

Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today	✓	✓	✓	✓	✓	✓	✓

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness	1	1	1	2	2	1	1
Irritability	0	0	1	0	0	1	1
Feelings of isolation	1	0	0	0	0	1	

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

Taking PRISTIQ

- Please mark or check each day of the week that you take your PRISTIQ

Symptom Tracking

Using the rating scale, describe how often you have experienced each of the depression symptoms on the list. Be sure to call a health care professional right away if you or a family member is experiencing recurrent thoughts of death or suicide.

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

- If your symptoms affect your ability to function on a daily basis, please note this in the comment box at the right-hand side of the Progress Tracker.

PRISTIQ Progress Tracker **Week 1**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

How to Take PRISTIQ

- Take PRISTIQ at about the same time each day.
- PRISTIQ may be taken with or without food.
- Swallow PRISTIQ tablets whole, with water.
- Do not divide, crush, chew, or dissolve PRISTIQ tablets because the tablets are time released.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 2**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

It may take several weeks before you start to feel better. Don't be discouraged, and don't stop taking PRISTIQ without talking to your health care professional, even if you do not feel results right away.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 3**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

You're not alone. Millions of people experience depression each year.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 4**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

Depression is not a sign of personal weakness or a condition that can be willed or wished away. You cannot merely “pull yourself together” and get better. But you may be successfully treated with prescription medication and other treatment options.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 5**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

Remember, you are not alone. Family members and friends, as well as your health care professional, may be able to help.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 6**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

Everyone responds differently to treatment. It's important to know that it could take several weeks to feel the full benefits of treatment. Remember to keep taking PRISTIQ exactly as prescribed by your health care professional.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 7**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

As treatment begins to take effect, you may notice some of the negative thoughts and feelings associated with depression fading.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

PRISTIQ Progress Tracker **Week 8**



Taking PRISTIQ	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Please mark or check each day of the week that you take your PRISTIQ.							
I took PRISTIQ today							

Symptom Tracking	Sun	Mon	Tue	Wed	Thu	Fri	Sat
Using the rating scale, describe how often you have experienced each of the following symptoms of depression this week.							
Constant sadness							
Irritability							
Feelings of isolation							
Muscle aches							
Low energy/fatigue							
Trouble sleeping							
Difficulty concentrating							
Lack of motivation							
Loss of interest in favorite activities							

Date:

Rating Scale

- 0 = Never
- 1 = Sometimes
- 2 = Most of the time

Bring this PRISTIQ Progress Tracker to your next appointment with your health care professional.

Do your symptoms affect your ability to function on a daily basis?

Message of Support

The fact is, depression is a real medical condition that cannot be willed or wished away. But it may be successfully treated with a prescription medication such as PRISTIQ.

Please note: This is not a formal diagnostic tool. Only a health care professional can actually diagnose depression. This may help you prepare for a discussion with your health care professional about depression symptoms you may have.

Please see the [Important Safety Information](#) on page 10 and the [Medication Guide for patients](#).

Important Safety Information

PRISTIQ™ (desvenlafaxine) is approved for the treatment of major depressive disorder in adults.



Suicidality and Antidepressant Drugs

Antidepressants increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. PRISTIQ is not approved for use in children under 18.

- People taking MAOIs should not take PRISTIQ.
- All patients taking antidepressants should be observed closely for signs that their condition is getting worse or that they are becoming suicidal. This is very important when an antidepressant is started or when the dose is changed. Patients should be watched for becoming agitated, irritable, hostile, aggressive, impulsive, or restless. These symptoms should be reported to the patient's health care professional right away.
- Tell your health care professional about all prescription and over-the-counter medications you are taking or plan to take, including:
 - Medicines to treat migraines or mood disorders, to avoid a potentially life-threatening condition
 - Aspirin, NSAID pain relievers, or blood thinners because they may increase the risk of bleeding
- PRISTIQ may cause or make some conditions worse, so tell your health care professional about all your medical conditions, including if you:
 - Have high blood pressure. Your blood pressure should be controlled before you start taking PRISTIQ and monitored regularly
 - Have heart problems, high cholesterol or triglyceride levels, or a history of stroke
 - Have glaucoma or increased eye pressure
 - Have kidney or liver problems
 - Have or had mania, bipolar disorder, or seizures or convulsions
 - Have low sodium levels in your blood
 - Are nursing, pregnant, or plan to become pregnant
- Discontinuation symptoms may occur when stopping PRISTIQ, especially when therapy is stopped suddenly. Talk to your health care professional before you stop taking or reduce the dose of PRISTIQ.
- Until you see how PRISTIQ affects you, be careful driving a car or operating machinery. Avoid drinking alcohol while taking PRISTIQ.
- Side effects when taking PRISTIQ 50 mg may include nausea, dizziness, sweating, constipation, and decreased appetite.

See the full [Medication Guide for patients](#) on page 11.

Medication Guide

PRISTIQ™ (pris-TEEK) Extended-Release Tablets (desvenlafaxine)



Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions

Read the Medication Guide that comes with you or your family member's antidepressant medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines. Talk to your, or your family member's, healthcare provider about:

- all risks and benefits of treatment with antidepressant medicines
- all treatment choices for depression or other serious mental illness

What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions?

1. Antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults within the first few months of treatment.
2. Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions. These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.
3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?
 - Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
 - Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
 - Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:

-
- | | |
|-------------------------------------|---|
| • thoughts about suicide or dying | • trouble sleeping (insomnia) |
| • attempts to commit suicide | • new or worse irritability |
| • new or worse depression | • acting aggressive, being angry, or violent |
| • new or worse anxiety | • acting on dangerous impulses |
| • feeling very agitated or restless | • an extreme increase in activity and talking (mania) |
| • panic attacks | • other unusual changes in behavior or mood |
-

Medication Guide (con't.)

What else do I need to know about antidepressant medicines?

- Never stop an antidepressant medicine without first talking to a healthcare provider. Stopping an antidepressant medicine suddenly can cause other symptoms.
- Antidepressants are medicines used to treat depression and other illnesses. It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- Antidepressant medicines have other side effects. Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- Antidepressant medicines can interact with other medicines. Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.
- Not all antidepressant medicines prescribed for children are FDA approved for use in children. Talk to your child's healthcare provider for more information.

This Medication Guide has been approved by the U.S. Food and Drug Administration for all antidepressants.

Important Information about Pristiq

Read the patient information that comes with Pristiq before you take Pristiq and each time you refill your prescription. There may be new information. If you have questions, ask your healthcare provider. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is Pristiq?

- Pristiq is a prescription medicine used to treat depression. Pristiq belongs to a class of medicines known as SNRIs (or serotonin-norepinephrine reuptake inhibitors).
- Pristiq has not been studied or approved for use in children and adolescents.

Who should not take Pristiq?

Do not take Pristiq if you:

- are allergic to desvenlafaxine, venlafaxine or any of the ingredients in Pristiq. See the end of this Medication Guide for a complete list of ingredients in Pristiq.
- currently take or have taken within the last 14 days, any medicine known as an MAOI. Taking an MAOI with certain other medicines, including Pristiq, can cause serious or even life-threatening side effects. Also, you must wait at least 7 days after you stop taking Pristiq before you take any MAOI.

Medication Guide (con't.)

What should I tell my healthcare provider before taking Pristiq?

Tell your healthcare provider about all your medical conditions, including if you:

- have high blood pressure
- have heart problems
- have high cholesterol or high triglycerides
- have a history of a stroke
- have glaucoma
- have kidney problems
- have liver problems
- have or had bleeding problems
- have or had seizures or convulsions
- have mania or bipolar disorder
- have low sodium levels in your blood
- are pregnant or plan to become pregnant. It is not known if Pristiq will harm your unborn baby.
- are breastfeeding. Pristiq can pass into your breast milk and may harm your baby. Talk with your healthcare provider about the best way to feed your baby if you take Pristiq.

Serotonin syndrome

A rare, but potentially life-threatening, condition called serotonin syndrome can happen when medicines such as Pristiq are taken with certain other medicines. Serotonin syndrome can cause serious changes in how your brain, muscles and digestive system work. Especially tell your healthcare provider if you take the following:

- medicines to treat migraine headaches known as triptans
- medicines used to treat mood disorders, including tricyclics, lithium, selective serotonin reuptake inhibitors (SSRIs), or serotonin norepinephrine reuptake inhibitors (SNRIs)
- silbutramine
- tramadol
- St. John's Wort
- MAOIs (including linezolid, an antibiotic)
- tryptophan supplements

Ask your healthcare provider if you are not sure if you are taking any of these medicines.

Before you take Pristiq with any of these medicines, talk to your healthcare provider about serotonin syndrome.

See *“What are the possible side effects of Pristiq?”*

Pristiq contains the medicine desvenlafaxine. Do not take Pristiq with other medicines containing venlafaxine or desvenlafaxine.

Medication Guide (con't.)

How should I take Pristiq?

- Take Pristiq exactly as your healthcare provider has told you.
- Take Pristiq at about the same time each day.
- Pristiq may be taken either with or without food.
- Swallow Pristiq tablets whole, with fluid. Do not crush, cut, chew, or dissolve Pristiq tablets because the tablets are time-released.
- When you take Pristiq, you may see something in your stool that looks like a tablet. This is the empty shell from the tablet after the medicine has been absorbed by your body.
- It is common for antidepressant medicines such as Pristiq to take several weeks before you start to feel better. Do not stop taking Pristiq if you do not feel results right away.
- Do not stop taking or change the dose of Pristiq without talking with your healthcare provider, even if you feel better.
- Talk with your healthcare provider about how long you should use Pristiq. Take Pristiq for as long as your healthcare provider tells you to.
- If you miss a dose of Pristiq, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose. Do not try to “make up” for the missed dose by taking two doses at the same time.
- Do not take more Pristiq than prescribed by your healthcare provider. If you take more Pristiq than the amount prescribed, contact your healthcare provider right away.
- In case of an overdose of Pristiq, call your healthcare provider or poison control center, or go to the emergency room right away.

What should I avoid while taking Pristiq?

- Do not drive a car or operate machinery until you know how Pristiq affects you.
- Avoid drinking alcohol while taking Pristiq.

What are the possible side effects of Pristiq?

Pristiq can cause serious side effects, including:

- See the beginning of this Medication Guide - Antidepressant Medicines, Depression and other Serious Mental Illnesses, and Suicidal Thoughts or Actions.
- Serotonin syndrome. See “What should I tell my healthcare provider before taking Pristiq?”

Get medical help right away if you think that you have serotonin syndrome. Signs and symptoms of serotonin syndrome may include one or more of the following:

-
- | | |
|--|------------------------------|
| • restlessness | • increase in blood pressure |
| • hallucinations (seeing and hearing things that are not real) | • diarrhea |
| • loss of coordination | • coma |
| • fast heart beat | • nausea |
| • increased body temperature | • vomiting |
-

Medication Guide (con't.)

Pristiq may also cause other serious side effects, including:

- New or worsened high blood pressure (hypertension). Your healthcare provider should monitor your blood pressure before and while you are taking Pristiq. If you have high blood pressure, it should be controlled before you start taking Pristiq.
- Abnormal bleeding or bruising. Pristiq and other SNRIs/SSRIs may cause you to have an increased chance of bleeding. Taking aspirin, NSAIDs (non-steroidal antiinflammatory drugs), or blood thinners may add to this risk. Tell your healthcare provider right away about any unusual bleeding or bruising.
- Glaucoma (increased eye pressure)
- Increased cholesterol and triglyceride levels in your blood
- Symptoms when stopping Pristiq (discontinuation symptoms). Side effects may occur when stopping Pristiq (discontinuation symptoms), especially when therapy is stopped suddenly. Your healthcare provider may want to decrease your dose slowly to help avoid side effects. Some of these side effects may include:

-
- | | |
|--------------------------------|-------------------|
| • dizziness | • anxiety |
| • nausea | • abnormal dreams |
| • headache | • tiredness |
| • irritability | • sweating |
| • sleeping problems (insomnia) | • diarrhea |
-

- Seizures (convulsions)
- Low sodium levels in your blood. Symptoms of this may include: headache, difficulty concentrating, memory changes, confusion, weakness and unsteadiness on your feet. In severe or more sudden cases, symptoms can include: hallucinations (seeing or hearing things that are not real), fainting, seizures and coma. If not treated, severe low sodium levels could be fatal.

Contact your healthcare provider if you think you have any of these side effects.

Medication Guide (con't.)

Common side effects with Pristiq include:



-
- | | |
|--------------------|----------------------------------|
| • nausea | • tiredness |
| • headache | • diarrhea |
| • dry mouth | • vomiting |
| • sweating | • anxiety |
| • dizziness | • tremor |
| • insomnia | • dilated pupils |
| • constipation | • decreased sex drive |
| • loss of appetite | • delayed orgasm and ejaculation |
| • sleepiness | |
-

These are not all the possible side effects of Pristiq. Tell your healthcare provider about any side effect that bothers you or does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. For more information on these and other side effects associated with Pristiq, talk to your healthcare provider, visit our web site at www.pristiq.com or call our toll-free number at 1-888-Pristiq.

How should I store Pristiq?

- Store Pristiq at 68° to 77°F (20° to 25°C).
- Do not use Pristiq after the expiration date (EXP), which is on the container. The expiration date refers to the last day of that month.
- Keep Pristiq and all medicines out of the reach of children.

General Information about the safe and effective use of Pristiq

Medicines are sometimes used for conditions that are not mentioned in Medication Guides. Do not use Pristiq for a condition for which it was not prescribed. Do not give Pristiq to other people, even if they have the same symptoms that you have. It may harm them. This Medication Guide summarizes the most important information about Pristiq. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about Pristiq that is written for healthcare professionals. For more information, go to www.pristiq.com or call 1-888-Pristiq (774-7847).

Medication Guide (con't.)

What are the ingredients in Pristiq?

Active ingredient: desvenlafaxine

Inactive ingredients: hypromellose, microcrystalline cellulose, talc, magnesium stearate, a film coating which consists of sodium carboxymethylcellulose, maltodextrin, dextrose, titanium dioxide, stearic acid and iron oxide(s).

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Issued February 2008.

Contact Information

Please visit our web site at www.pristiq.com, or call our toll-free number 1-888-Pristiq to receive more information.



This product's label may have been updated. For current package insert and further product information, please visit www.wyeth.com or call our medical communications department toll-free at 1-800-934-5556.



Wyeth®

Wyeth Pharmaceuticals Inc.

Philadelphia, PA 19101

W10530P002

ET01

Rev 04/08